Applicants: David J. Pinsky, et al.

U.S. Serial No.: 10/679,135

Filed: October 3, 2003

Page 2

Amendments to the Specification

Please replace the paragraph beginning on page 1, at line 5 of the specification, a paragraph which was added to the specification by a Preliminary Amendment filed on October 3, 2003, with the following:

This application is a continuation continuation-inpart of U.S. Serial No. 09/671,100, filed September
27, 2000, now abandoned, which is a continuation of
PCT International Application No. PCT/US99/07175,
filed April 1, 1999., which is a continuation inpart of U.S. Serial No. 09/053,871, filed April 1,
1998, now U.S. Patent No. 6,315,995 B1, issued
November 13, 2001 which is a continuation in part of
PCT International Application No. PCT/US97/17229,
filed September 25, 1997, and a continuation in part
of U.S. Serial No. 08/721,447, filed September 27,
1996, all of which applications are hereby
incorporated by reference in their entireties into
this application.

Please replace the sentence in the section starting on page 3, line 1 with the following rewritten sentence:

Brief Description of the Figures Drawings

Please replace the paragraph in the "Brief Description of the Drawing" section on page 9, line 30 to page 10, line 6 for Figure 19, with the following rewritten paragraph:

Figure 19 19A, 19B and 19C: Role of P-selectin in the cerebrovascular no-reflow. Cerebral blood flow was measured in PS +/+ (Fig 19A top panel) and PS -/- (Fig 19B middle panel) mice using a laser doppler flow probe, and expressed as the percentage

Applicants: David J. Pinsky, et al.

U.S. Serial No.: 10/679,135

Filed: October 3, 2003

Page 3

of contralateral (nonischemic) hemispheric blood flow (\pm SEM). Blood flow was measured at the following time points: **a**, prior to MCAO (PS +/+, n = 16; PS -/-, n = 23); **b**, immediately following MCAO (PS +/+, n = 42; PS -/-, n = 40); **c**, 10 minutes following MCAO but still prior to reperfusion (PS +/+, n = 36; PS -/-, n = 34); **d**, immediately following reperfusion (PS +/+, n = 36; PS -/-, n = 34); **e**, 30 minutes following reperfusion (PS +/+, n = 8; PS -/-, n = 5); and **f**, 22 hours following reperfusion (PS +/+, n = 15; PS -/-, n = 5). The bottom panel Fig 19C represents an overlay of the top two panels, with error bars omitted for clarity.